

REFERENCES

- Agre, P., Preston, G. M., Smith, B. L., Jung, J. S., Raina, S., Moon, C., Guggino, W. B., & Nielsen, S. (1993a). Aquaporin CHIP: the archetypal molecular water channel. *American Journal of Physiology – Renal Physiology* **265**, F463–F476.
- Agre, P., Sasaki, S., & Chrispeels J. (1993b). Aquaporins: a family of water channel proteins. *American Journal of Physiology – Renal Physiology* **265**, F461.
- Agre, P., King, L. S., Yasui, M., Guggino, W. B., Ottersen, O. P., Fujiyoshi, Y., Engel, A., and Nielsen, S. (2002). Aquaporin water channels – From atomic structure to clinical medicine. *Journal of Physiology* **542**, 3–16.
- Amiry-Moghaddam, M., Otsuka, T., Hurn, P. D., Traystman, R. J., Finn-Mogens, H., Froehner, S. C., Adams, M. E., Neely, J. D., Agre, P., Ottersen, O. P., Bhardwaj, A. (2003) An alpha-syntrophin-dependent pool of AQP4 in astroglial end-feet confers bidirectional water flow between blood and brain. *Proceedings of the National Academy of Sciences of the USA* **100**, 2106–2111.
- Beroukas, D., Hiscock, J., Gannon, B. J., Jonsson, R., Gordon, T. P., & Waterman, S. A. (2002). Selective down-regulation of aquaporin-1 in salivary glands in primary Sjögren's syndrome. *Laboratory Investigation* **82**, 1547–1552.
- Berry, V., Francis, P., Kaushal, S., Moore, A., & Bhattacharya, S. (2000). Missense mutations in MIP underlie autosomal dominant 'polymorphic' and lamellar cataracts linked to 12q. *Nature Genetics* **25**, 15–17.
- Calamita, G., Bishai, W. R., Preston, G. M., Guggino, W. B., & Agre, P. (1995). Molecular cloning and characterization of AqpZ, a water channel from *Escherichia coli*. *Journal of Biological Chemistry* **270**, 29063–29066.
- Carbrey, J. M., Gorelick-Feldman, D. A., Kozono, D., Praetorius, J., Nielsen, S., & Agre, P. (2003). Aquaglyceroporin AQP9: Solute permeation and metabolic control of expression in liver. *Proceedings of the National Academy of Sciences of the USA* **100**, 2945–2950.
- Chou, C. L., Knepper, M. A., van Hoek, A. N., Brown, D., Yang, B., Ma, T., & Verkman, A. S. (1999). Reduced water permeability and altered ultrastructure in thin descending limb of Henle in aquaporin-1 null mice. *Journal of Clinical Investigation* **103**, 491–496.
- de Groot, B. L. & Grubmuller, H. (2001). Water permeation across biological membranes: mechanism and dynamics of aquaporin-1 and GlpF. *Science* **294**, 2353–2357.
- Deen, P. M., Verdijk, M. A., Knoers, N. V., Wieringa, B., Monnens, L. A., van Os, C. H., & Van Oost, B. A. (1994). Requirement of human renal water channel aquaporin-2 for vasopressin-dependent concentration of urine. *Science* **264**, 92–95.
- Denker, B. M., Smith, B. L., Kuhajda, F. P., & Agre, P. (1988). Identification, purification, and partial characterization of a novel Mr 28,000 integral membrane protein from erythrocytes and renal tubules. *Journal of Biological Chemistry* **263**, 15634–15642.
- Finkelstein, A. (1987). Water movement through lipid bilayers, pores, and plasma membranes: Theory and reality. New York: Wiley.
- Francis, P., Chung, J. J., Yasui, M., Berry, V., Moore, A., Wyatt, M. K., Wistow, G., Bhattacharya, S. S., & Agre, P. (2000). Functional impairment of lens aquaporin in two families with dominantly inherited cataracts. *Human Molecular Genetics* **9**, 2329–2334.
- Frigeri, A., Nicchia, G. P., Nico, B., Quondamatteo, F., Herken, R., Roncali, L., & Svelto, M. (2001). Aquaporin-4 deficiency in skeletal muscle and brain of dystrophic mdx mice. *FASEB Journal*. **15**, 90–98.
- Fu, D., Libson, A., Miercke, L. J., Weitzman, C., Nollert, P., Krucinski, J., & Stroud, R. M. (2000). Structure of a glycerol-conducting channel and the basis for its selectivity. *Science* **290**, 481–486.
- Furman, C. S., Gorelick-Feldman, D. A., Davidson, K. G. V., Yasumura, T., Neely, J. D., Agre, P., & Rash, J. E. (2003). Aquaporin-4 square array assembly: opposing actions of M1 and M23 isoforms. *Proceedings of the National Academy of Sciences of the USA* **100**, 13609–13614.
- Fushimi, K., Uchida, S., Hara, Y., Hirata, Y., Marumo, F., & Sasaki, S. (1993). Cloning and expression of apical membrane water channel of rat kidney collecting tubule. *Nature* **361**, 549–552.
- Hohmann, S. (2002). Osmotic stress signaling and osmoadaptation in yeasts. *Microbiology and Molecular Biology Reviews* **66**, 300–372.

- Jung, J. S., Bhat, R. V., Preston, G. M., Guggino, W. B., Baraban, J. M., & Agre, P. (1994a). Molecular characterization of an aquaporin cDNA from brain: candidate osmoreceptor and regulator of water balance. *Proceedings of the National Academy of Sciences of the USA* **91**, 13052–13056.
- Jung, J. S., Preston, G. M., Smith, B. L., Guggino, W. B., & Agre, P. (1994b). Molecular structure of the water channel through aquaporin CHIP. The hourglass model. *Journal of Biological Chemistry* **269**, 14648–14654.
- Kaldenhoff, R., Grote, K., Zhu, J. J., Zimmermann, U. (1998). Significance of plasma lemma aquaporins for water-transport in *Arabidopsis thaliana*. *Plant Journal* **14**, 121–128.
- King, L. S., Nielsen, S., & Agre, P. (1996). Aquaporin-1 water channel protein in lung: ontogeny, steroid-induced expression, and distribution in rat. *Journal of Clinical Investigation* **97**, 2183–2191.
- King, L. S., Nielsen, S., & Agre, P. (1997). Aquaporins in complex tissues. I. Developmental patterns in respiratory and glandular tissues of rat. *American Journal of Physiology – Cell Physiology* **273**, C1541–C1548.
- King, L. S., Choi, M., Fernandez, P. C., Cartron, J. P., & Agre, P. (2001). Defective urinaryconcentrating ability due to a complete deficiency of aquaporin-1. *New England Journal of Medicine* **345**, 175–179.
- King, L. S., Nielsen, S., Agre, P., & Brown, R. H. (2002). Decreased pulmonary vascular permeability in aquaporin-1-null humans. *Proceedings of the National Academy of Sciences of the USA* **99**, 1059–1063.
- Kozono, D., Yasui, M., King, L. S., & Agre, P. (2002). Aquaporin water channels: atomic structure and molecular dynamics meet clinical medicine. *Journal of Clinical Investigation* **109**, 1395–1399.
- Kozono, D., Ding, X., Iwasaki, I., Meng, X., Kamagata, Y., Agre, P., & Kitagawa, Y. (2003). Functional expression and characterization of an archaeal aquaporin: AqpM from *Methanothermobacter marburgensis*. *Journal of Biological Chemistry* **278**, 10649–10656.
- Kuriyama, H., Shimomura, H., Kishida, K., Kondo, H., Furuyama, N., Nishizawa, H., Maeda, N., Matsuda, M., Nagaretani, H., Kihara, S., Nakamura, T., Tochino, Y., Funahashi, T., Matsuzawa, Y. (2002). Coordinated regulation of fat-specific and liver-specific glycerol channels, aquaporin adipose, and aquaporin 9. *Diabetes* **51**, 2915–2921.
- Liu, Z., Shen, J., Carbrey, J.M., Mukhopadhyay, R., Agre, P., & Rosen, B. (2002). Arsenite transport by mammalian aquaglyceroporins AQP7 and AQP9. *Proceedings of the National Academy of Sciences of the USA* **99**, 6053–6058.
- Macey, R. I. & Farmer, R. E. L. (1970) Inhibition of water and solute permeability in human red cells. *Biochimica Biophysica Acta* **211**, 104–106.
- Manley, G. T., Fujimura, M., Ma, T., Noshita, N., Filiz, F., Bollen, A. W., Chan, P., & Verkman, A. S. (2000). Aquaporin-4 deletion in mice reduces brain edema after acute water intoxication and ischemic stroke. *Nature Medicine* **6**, 159–163.
- Moon, C., Preston, G. M., Griffin, C. A., Jabs, E. W., & Agre, P. (1993). The human aquaporin-CHIP gene. Structure, organization, and chromosomal localization. *Journal of Biological Chemistry* **268**, 15772–15778.
- Murata, K., Mitsuoka, K., Hirai, T., Walz, T., Agre, P., Heymann, J. B., Engel, A., & Fujiyoshi, Y. (2000). Structural determinants of water permeation through aquaporin-1. *Nature* **407**, 599–605.
- Nagelhus, E. A., Veruki, M. L., Torp, R., Haug, F. M., Laake, J. H., Nielsen, S., Agre, P., & Ottersen, O. P. (1998). Aquaporin-4 water channel protein in the rat retina and optic nerve: polarized expression in Muller cells and fibrous astrocytes. *Journal of Neuroscience* **18**, 2506–2519.
- Nakhoul, N. L., Davis, B. A., Romero, M. F., & Boron, W. F. (1998). Effect of expressing the water channel aquaporin-1 on the CO₂ permeability of *Xenopus* oocytes. *American Journal of Physiology – Cell Physiology* **274**, C297–298.
- Neely, J. D., Amiry-Moghaddam, M., Ottersen, O. P., Froehner, S. C., Agre, P., & Adams, M. E. (2001). Syntrophin-dependent expression and localization of Aquaporin-4 water channel protein. *Proceedings of the National Academy of Sciences of the USA* **98**, 14108–14113.

Nejsum, L. N., Kwon, T. H., Jensen, U. B., Fumagalli, O., Frokiaer, J., Krane, C. M., Menon, A. G., King, L. S., Agre, P., & Nielsen, S. (2002). Functional requirement of aquaporin-5 in plasma membranes of sweat glands. *Proceedings of the National Academy of Sciences of the USA* **99**, 511–516.

Nielsen, S., DiGiovanni, S.R., Christensen, E.I., Knepper, M.A., & Harris, H.W. (1993a). Cellular and subcellular immunolocalization of vasopressin-regulated water channel in rat kidney. *Proceedings of the National Academy of Sciences of the USA* **90**, 11663–11667.

Nielsen, S., Smith, B. L., Christensen, E. I., & Agre, P. (1993b). Distribution of the aquaporin CHIP in secretory and resorptive epithelia and capillary endothelia. *Proceedings of the National Academy of Sciences of the USA* **90**, 7275–7279.

Nielsen, S., Smith, B. L., Christensen, E. I., Knepper, M. A., & Agre, P. (1993c). CHIP28 water channels are localized in constitutively water-permeable segments of the nephron. *Journal of Cell Biology* **120**, 371–383.

Nielsen, S. & Agre, P. (1995). The aquaporin family of water channels in kidney. *Kidney International* **48**, 1057–1068.

Nielsen, S., Chou, C. L., Marples, D., Christensen, E. I., Kishore, B. K., & Knepper, M. A. (1995). Vasopressin increases water permeability of kidney collecting duct by inducing translocation of aquaporin-CD water channels to plasma membrane. *Proceedings of the National Academy of Sciences of the USA* **92**, 1013–1017.

Nielsen, S., Nagelhus, E. A., Amiry-Moghaddam, M., Bourque, C., Agre, P., & Ottersen, O. P. (1997). Specialized membrane domains for water transport in glial cells: high-resolution immunogold cytochemistry of aquaporin-4 in rat brain. *Journal of Neuroscience* **17**, 171–180.

Nielsen, S., Frokiaer, J., Marples, D., Kwon, T. H., Agre, P., & Knepper, M. A. (2002). Aquaporins in the kidney: from molecules to medicine. *Physiological Reviews* **82**, 205–244.

Preston, G. M. & Agre, P. (1991). Isolation of the cDNA for erythrocyte integral membrane protein of 28 kilodaltons: member of an ancient channel family. *Proceedings of the National Academy of Sciences of the USA* **88**, 11110–11114.

Preston, G. M., Carroll, T. P., Guggino, W. B., & Agre, P. (1992). Appearance of water channels in *Xenopus* oocytes expressing red cell CHIP28 protein. *Science* **256**, 385–387.

Preston, G. M., Jung, J. S., Guggino, W. B., & Agre, P. (1993). The mercury-sensitive residue at cysteine 189 in the CHIP28 water channel. *Journal of Biological Chemistry* **268**, 17–20.

Preston, G. M., Smith, B. L., Zeidel, M. L., Moulds, J. J., & Agre, P. (1994). Mutations in aquaporin-1 in phenotypically normal humans without functional CHIP water channels. *Science* **265**, 1585–1587.

Raina, S., Preston, G. M., Guggino, W. B., & Agre, P. (1995). Molecular cloning and characterization of an aquaporin cDNA from salivary, lacrimal, and respiratory tissues. *Journal of Biological Chemistry* **270**, 1908–1912.

Ramon y Cajal, S. (1897). *Advice to a Young Investigator* (translated by N. Swanson and L. W. Swanson, 1999) Cambridge, MIT Press.

Schnermann, J., Chou, C. L., Ma, T., Traynor, T., Knepper, M. A., & Verkman, A. S. (1998). Defective proximal tubular fluid reabsorption in transgenic aquaporin-1 null mice. *Proceedings of the National Academy of Sciences of the USA* **95**, 9660–9664.

Smith, B. L. & Agre, P. (1991). Erythrocyte Mr 28,000 transmembrane protein exists as a multisubunit oligomer similar to channel proteins. *Journal of Biological Chemistry* **266**, 6407–6415.

Smith, B. L., Preston, G. M., Spring, F. A., Anstee, D. J., & Agre, P. (1994). Human red cell aquaporin CHIP. I. Molecular characterization of ABH and Colton blood group antigens. *Journal of Clinical Investigation* **94**, 1043–1049.

Steinfeld, S., Cogan, E., King, L. S., Agre, P., Kiss, R., & Delporte, C. (2001). Abnormal distribution of aquaporin-5 water channel protein in salivary glands from Sjögren's syndrome patients. *Laboratory Investigation* **81**, 143–148.

Sui, H., Han, B. G., Lee, J. K., Walian, P., & Jap, B. K. (2001). Structural basis of water-specific transport through the AQP1 water channel. *Nature* **414**, 872–878.

Tajkhorshid, E., Nollert, P., Jensen, M. O., Miercke, L. J., O'Connell, H., Stroud, R. M., &

Schulten, K., (2002). Control of the selectivity of the aquaporin water channel family by global orientational tuning. *Science* **296**, 525–530.

Tournaire-Roux, C., Sutka, M., Javot, H., Gout, E., Gerbeau, P., Luu, D. T., Bligny, R. & Maurel, C. (2003). Cytosolic pH regulates root water transport during anoxic stress through gating of aquaporins. *Nature* **425**, 393–397.

Tsubota, K., Hirai, S., King, L. S., Agre, P., & Ishida, N. (2001). Defective cellular trafficking of lacrimal gland aquaporin-5 in Sjögren's syndrome. *Lancet* **357**, 688–689.

Uehlein, N., Lovisolo, C., Siefritz, F., & Kaldenhoff, R. (2003). The tobacco aquaporin NtAQP1 is a membrane CO₂ pore with physiological functions. *Nature* **425**, 734–737.

Vajda, Z., Pedersen, M., Fuchtbauer, E.-M., Wert, K., Stodkilde-Jorgensen, H., Sulyok, E., Doczi, T., Neely, J. D., Agre, P., Frokiaer, J., & Nielsen, S. (2002). Delayed onset of brain edema and mislocalization of aquaporin-4 in dystrophin-null transgenic mice. *Proceedings of the National Academy of Sciences of the USA* **99**, 13131–13136.

Walz, T., Smith, B. L., Zeidel, M. L., Engel, A., & Agre, P. (1994). Biologically active two-dimensional crystals of aquaporin CHIP. *Journal of Biological Chemistry* **269**, 1583–1586.

Yasui, M., Hazama, A., Kwon, T. H., Nielsen, S., Guggino, W. B., & Agre, P. (1999). Rapid gating and anion permeability of an intracellular aquaporin. *Nature* **402**, 184–187.

Zeidel, M. L., Ambudkar, S. V., Smith, B. L., & Agre, P. (1992). Reconstitution of functional water channels in liposomes containing purified red cell CHIP28 protein. *Biochemistry* **31**, 7436–7440.

Zeidel, M. L., Nielsen, S., Smith, B. L., Ambudkar, S. V., Maunsbach, A. B., & Agre, P. (1994). Ultrastructure, pharmacologic inhibition, and transport selectivity of aquaporin channel-forming integral protein in proteoliposomes. *Biochemistry* **33**, 1606–1615.

Zeuthen, T. & Klaerke, D. A. (1999). Transport of water and glycerol in aquaporin 3 is gated by H(+). *Journal of Biological Chemistry* **274**, 21631–21636.